

*Rhyparus helepholoides* Fairmaire, 1893 ヒメセスジカクマグソコガネ

*Rhyparus helepholoides* Fairmaire, 1893, Note Lyden Mus. XV, p.145. (Borneo)

*Rhyparus amamianus* Nakane, 1956, Ins. Mats., Sapporo 20(3/4):123-124. (Amami-Oshima)

Distr. Japan (Tokara Is., - Suwanose-jima, Takara-jima, Nakanoshima, Amami-Oshima, Okinoerabu-jima, Okinawa-honto, Ishigaki-jima, Iriomote-jima, Miyako-jima), Taiwan, Borneo.

*Rhyparus kitanoi* Y. Miyake, 1982 キュウシュウカクマグソコガネ

*Rhyparus kitanoi* Y. Miyake, 1982, Kita Kyushu no Konchu 29(2):65-68, fig.1-8. (Cape Sata, Ohsumi, Meshima-Danjo Is., Akamatsu Valley, Shimabara City)

Distr. Japan (Kyushu-Mainland and Danjo Is.).

*Rhyparus peninsularis* Arrow, 1905 セスジカクマグソコガネ

*Rhyparus peninsularis* Arrow, 1905, Ann. Mag. Nat. Hist. (7)XV, p.537.

*Rhyparus azumai* Nakane, 1956, Ins. Mats., Sapporo 20(3/4):122-123, fig. (Kii, Hiko, Chikugo)

Distr. Japan (Honshu, Nii-jima, Miyake, Mikura, Hachijyo-jima, Shikoku, Kyushu, Yakushima, Tanegashima, Nakano-shima, Takara-jima, Tokara Is., Okinawa, Goto Is., Danjo Is., Tsushima, Hekura-jima, Ishigaki-jima), Taiwan.

## Subfamily AEGIALIINAE ニセマグソコガネ亜科

Genus *Aegialia* Latrille, 1807 ニセマグソコガネ属

*Aegialis* Latreille, 1807, Gen. Crust. et Ins. II, 1807, p.96.

(type-species : *Scarabaeus arearia* Fabricius, 1787)

*Aegialia nitida* Waterhouse, 1875 ニセマグソコガネ

*Aegialia nitida* Waterhouse, 1875, Trans. ent. Soc. Lond., p.95. (Hakodate)

Distr. Japan (Hokkaido, Honshu, Kyushu).

Genus *Caelius* Lewis, 1895 トゲマグソコガネ属

*Caelius* Lewis, 1895, Ann. Mag. Nat. Hist. (6)XVI, p.386, f.5.

(type-species : *Caelius denticollis* Lewis, 1895)

*Caelius denticollis* Lewis, 1895 トゲマグソコガネ

*Caelius denticollis* Lewis, 1895, Ann. Mag. Nat. Hist. ser.6, 16, p.381. (Miyanosita, Kiga, Nikko)

Distr. Japan (Hokkaido, Honshu, Kyushu).

Genus *Psammoporus* Thomson, 1863 ナガニセマグソコガネ属

*Psammoporus* Thomson, 1863, Skand. Col. (5). p.72.

(type-species : *Scarabaeus sabuleti* Panzer, 1797)

*Psammoporus comis* (Lewis, 1895) ナガニセマグソコガネ

- Psammobius comis* Lewis, 1895, Ann. Mag. Nat. Hist. (6) 16:384.  
*Aegialia (Dimalia) hybrida* Reitter, 1892, Verh. Nat. Ver. Brun. XXX, 189:251.  
*Psammoporus freyi* Portrovitz, 1961, Arb. Mus. Frey. 12(1):136.  
 Distr. Japan (Hokkaido, Honshu).

- Psammoporus kamtschaticus* (Motschulsky, 1860) アラメニセマグソコガネ  
*Aegialia kamtschatica* Motschulsky, 1860, Dr. Schrenk's Reisen u Forsch im Amur-Lande II, 2, Lief.:132,  
 pl.9, f.3. (Kamtschatka & Daoure)  
*Psammoporus nakanei* Masumoto, 1986, Pap. Ent. Pres. Nakane, Tokyo pp.144-164, figs.1&4.  
 Distr. Japan (Hokkaido, Honshu), Kuriles, Kamtschatake, Sachalin, E.Siberia.

- Psammoporus tsukamotoi* Masumoto, 1986 キタアラメニセマグソコガネ  
*Psammoporus tsukamotoi* Masumoto, 1986, Pap. Ent. Pres. Nakane Tokyo p.146-147, figs.2&5. (Mt. Rishiri,  
 Rishiri Is., Horomitoge, Sounkyo, Mt. Poroshiridake, Hidaka-Hokkaido)  
 Distr. Japan (Hokkaido, Rishiri Is.).

**Subfamily GLAPHYRINAE ヒゲフトハナムグリ亜科**

- Genus *Amphicoma* Latreille, 1807 ヒゲフトハナムグリ属**  
*Amphicoma* Latreille, 1807, Genera Crust. et. Insect, 2:118.  
*Anthypna* Latreille, 1807, Genera Crust. et. Insect, 2:119.  
*Anthynoides* Yawata, 1942, Trans. Kansai Ent. Soc. 12(1):33-35.  
 (type-species : *Melolontha abdominalis* Fabricius, 1781)

- Amphicoma pectinata* (Lewis, 1895) ヒゲフトハナムグリ  
*Anthypna pectinata* Lewis, 1895, Ann. Mag. Nat. Hist. (6) XVI, p.358. (Tokio)  
 Distr. Japan (Honshu, Shikoku).

- Amphicoma splendens* (Yawata, 1942) オオヒゲフトハナムグリ  
*Anthypnoides splendens* Yawata, 1942, Trans. Kansai Ent. Soc. 12(1):35. (Yonakuni Is.)  
 Distr. Japan (Ishigaki Is., Iriomote Is., Yonaguni Is.).

**Subfamily MELOLONTHINAE コフキコガネ亜科**

**Tribe HOPLIINI アシナガコガネ族**

- Genus *Hoplia* Illiger, 1803 アシナガコガネ属**  
*Hoplia* Illiger, 1803, Mag. Ins. II, p.226.  
 (type-species : *Scarabaeus farinosus* Linne, 1761)  
*Hyperis* Dejean, 1833, Cat. Col. ed. 3, p.167.  
*Echyra* Erichson, 1847, Nat. Ins. Detsch. III, p.705, nota.