

Distr. Japan (Tsushima?), Quelpart Is., Korea, S. Manchuria, China.

Anomala testaceipes (Motschulsky, 1860) スジコガネ

Rhomobonyx testaceipes Motschulsky, 1860, Etud. Ent. IX, p.14. (Japan)

Mimela (Paramimela) testaceipes var. *ochrochalcea* Ohaus, 1915, Stett. ent. Ztg., 76, p.91.

Mimela (Paramimela) testaceipes var. *ochroptera* Ohaus, 1915, Stett. ent. Ztg., 76, p.91.

Mimela (Paramimela) testaceipennis var. *anocybria* Ohaus, 1915, Stett. ent. Ztg., 76, p.92.

Mimela (Paramimela) testaceipennis var. *cyprionoptera* Ohaus, 1915, Stett. ent. Ztg., 76, p.92.

Distr. Japan (Hokkaido, Honshu, Sado, Izu Is.—Oshima, Nii—jima, Toshima, Shikine—jima, Koze—jima, Shikoku, Kyushu, Goto, Hirato, Iki, Tsushima, Yakushima, Tokara), Saghalin, Korea, China, E. Siberia.

subsp. *yusai* Nomura, 1969

Anomala testaceipes yusai Nomura, 1969, Ent. Rev. Japan 21 (2):73. (Miyake—jima, Mikura—jima, Izu Is.)

Distr. Japan (Miyake—jima, Mikura—jima).

Anomala triangularis Schoenfeldt, 1890 サンカクスジコガネ

Anomala triangularis Schoenfeldt, 1890, Ent. Nachr. XVI, 11, p.171. (Oshima)

Distr. Japan.

subsp. *kyushuensis* Nomura, 1964 キュウシュウスジコガネ

Anomala triangularis kyushuensis Nomura, 1964, Tōhō Gakuho (14):68. (Ohsumi)

Distr. Japan (Kyushu, Hekura—jima).

subsp. *triangularis* Schoenfeldt, 1890 サンカクスジコガネ

Anomala triangularis Schoenfeldt, 1890, Ent. Nachr., XVI, II, p.171. ('shima)

Distr. Japan (Tanegashima, Tokara Is.—Akuseki—jima, Takara—jima, Nakano—shima, Amami—Oshima, Tokuno—shima, Okinoerabu—jima, Yoronoto, Okinawa).

Anomala xanthopleura Arrow, 1913 リュウキュウドウガネ

Anomala xanthopleura Arrow, 1913, Ann. Mag. Nat. Hist. (8)XII, p.402. (Loo—Choo)

Distr. Japan (Amami—Oshima, Okinawa Is. Ishigaki—jima, Kume—jima).

Subfamily DYNASTINAE カブトムシ亜科

Tribe DYNASTINI カブトムシ族

Genus *Allomyrina* Arrow, 1911 カブトムシ属

Allomyrina Arrow, 1911, Ann. Mag. Nat. Hist. (8)XII, p.153.

(type—species : *Scarabaeus dichotomus* Linne, 1771)

Allomyrina dichotoma (Linné, 1771) カブトムシ

Scarabaeus dichotomus Linne, 1771, Mant. Plant. VI, p.529.

Distr. Japan, Taiwan, Korea, China, Indochina.

subsp. *inchachina* Kusui, 1976 クメジマカブトムシ

Allomyrina dichotoma inchachina Kusui, 1976, Ent. Rev. Japan, 29 (1/2):52—53, pl.5. (Kume—jima)

- Distr. Japan (Kume-jima).
 subsp. *septentrionalis* (Kōno, 1931) カブトムシ
Trypoxylus dichotomus Linne, var. *septentrionalis* Kōno, 1931, Ins. Mats., V, p.160, t.3, f.5. (Honshu-Iwate)
Allomyrina dichotoma septentrionalis, Kurosawa, 1985, Coleop. Jap. Col. II, pl.76, f.1, p.417.
 Distr. Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Tanegashima, Yakushima, Kuchinoerabu-jima), Korea, N.China, Indo-china.
 subsp. *takarai* Kusui, 1976 オキナワカブトムシ
Allomyrina dichotoma takarai Kusui, 1976, Ent. Rev. Japan 29(1/2):52, pl.5. (Okinawa Is.)
 Distr. Japan (Okinawa Is.).

Tribe ORYCTINI サイカブトムシ族

- Genus *Oryctes* Illiger, 1798 サイカブトムシ属
Oryctes Illiger, 1798, Kafer Fransseus, p.11.
 (type-species : *Scarabaeus nasicornis* Linne, 1758)

- Oryctes rhinoceros* (Linne, 1758) サイカブトムシ
Scarabaeus rhinoceros Linne, 1758, Syst. Nat. p.346.
 Distr. Japan (Yonaguni Is., Ishigaki Is.), Korea, China, Taiwan, Indo-China, Siam, Burma, India, Malay, Philippine, Sumatora, Java, Celebes, Ceram, Amboyna.

- Genus *Eophileurus* Arrow, 1908 コカブトムシ属
Eophileurus Arrow, 1908, Trans. ent. Soc. London, p.332.
 (type-species : *Geotrupes planatus* Wiedemann, 1823)

- Eophileurus chinensis* (Faldermann, 1835) コカブトムシ
Phileurus chinensis Faldermann, 1835, Mem. Acad. Petr. ii, p.370.
Trionychus poteli Fairmaire, 1898, Ann. Soc. Ent. France, p.384.
 Distr. Japan (Hokkaido, Honshu, Tobishima, Sado, Izu Is.-Oshima, Toshima, Nii-jima, Kozu-jima, Miyake-jima, Mikura-jima, Shikoku, Kyushu), Saisyu-to, Taiwan, Burma, Ussurie, China).
 subsp. *irregularis* Prell, 1913 アマミコカブトムシ
Eophileurus chinensis var. *irregularis* Prell, 1913, Suppl. Ent. ii, p.23.
 Distr. Japan (Amami-Oshima).
 subsp. *okinawanus* Nomura, 1964 オキナワコカブトムシ
Eophileurus chinensis okinawanus Nomura, 1964, Ent. Rev. Japan 16(2):57, pl.8, f.12. (Okinawa Is., Iriomote Is.)
 Distr. Japan (Okinawa Is., Iriomote Is.).

Tribe PENTODONTINI クロマルコガネ族

Genus *Alissonotum* Arrow, 1908 クロマルコガネ属

Alissonotum Arrow, 1908, Trans. ent. Soc. London, p.332.

(type-species : *Geotrupes piceus* Fabricius, 1775)

Alissonotum pauper (Burmeister, 1847) クロマルコガネ

Heteronychus pauper Burmeister, 1847, Handb. Ent. V, p.94. (Philippines)

Distr. Japan (Okinoerabu Is., Tokara Is.), Taiwan, China, Philippines, Indo-China.

Subfamily EUCHIRINAE テナガコガネ亜科

Genus *Cheirotonus* Hope, 1841 テナガコガネ属

Cheirotonus Hope, 1841, Trans. Linn. Soc. (17), p.594.

(type-species : *Cheirotonus macleayi* Hope, 1841)

Cheirotonus jambar Y. Kurosawa, 1984 ヤンバルテナガコガネ

Cheirotonus jambar Y. Kurosawa, 1984, Bull. Natn. Sci. Mus. Tokyo, Ser. A, 10(2):73-78, fig.1-3. (N.Okinawa Is.)

Distr. Japan (N.Okinawa Is.).

Subfamily CETONIINAE ハナムグリ亜科

Tribe HETERORRHININI カナブン族

Genus *Cosmiomorpha* Saunders, 1852 チャイロカナブン属

Cosmiomorpha Saunders, 1852, Trans. ent. Soc. London (n.s.) II, p.28.

(type-species : *Cosmiomorpha modesta* Saunders, 1852)

Subgenus *Microcosmiomorpha* Mikšić, 1974

Microcosmiomorpha Mikšić, 1974, Rev. suis. Zool. LXXXI, 4, p.770.

(type-species : *Cosmiomorpha setulosa* Westwood, 1854)

Cosmiomorpha (Microcosmiomorpha) similis Fairmaire, 1899 チャイロカナブン

Cosmiomorpha similis Fairmaire, 1899, Ann. Soc. Ent. Fr. LXVII, p.621.

Distr. Japan, Taiwan.

subsp. *nigra* Nijima et Kinoshita, 1927 チャイロカナブン

Cosmiomorpha similis Fairmaire var. *nigra* Nijima et Kinoshita, 1927, Res. Bull. Coll. Exp. Forest Coll. Agr. Hokkaido Imp. Univ. Japan IV:69. (Ishigaki-jima)

Distr. Japan (Ishigaki-jima, Iriomote-jima, Obama-jima).

subsp. *miyakoana* Nomura, 1964 ミヤコチャイロカナブン

Cosmiomorpha similis miyakoana Nomura, 1964, Tōhō Gakuho (14):70. (Miyako Is.)

Distr. Japan (Miyako-jima, Irabu-jima).